

**NEW EVIDENCE FOR ALTERNATING EFFUSIVE AND EXPLOSIVE
ERUPTIONS FROM THE TYPE SECTION OF THE STANISLAUS GROUP IN
THE “CATARACT” PALEOCANYON, CENTRAL SIERRA NEVADA**

Koerner, Alice, and Busby, Cathy, Department of Earth Science, University of
California, Santa Barbara, CA 93106, USA

Putirka, Keith, and Pluhar, Christopher J., Department of Earth and Environmental
Sciences, California State University-Fresno, 2345 San Ramon Ave., MS/MH24,
Fresno, CA 93720, USA

ABSTRACT

High-K volcanic rocks of the Stanislaus Group record magmatic events that occurred when the Sierra Nevada microplate began to calve off the western edge of the Nevada Plano at 10–9 Ma. Despite the fact that these rocks have been studied in the Sierra Nevada for 120 years, their stratigraphy is not well known, because they were largely deposited in a complicated paleocanyon network that drained the Nevada Plano and are preserved on high peaks in roadless areas. We present new 1:6,000 scale mapping and a detailed measured section in the Bald Peak-Red Peak area of Sonora Pass, in the upper reach of the “Cataract” paleocanyon, or paleo-Stanislaus River. The Bald Peak-Red Peak area was proposed as the type section for the Stanislaus Group in 1953 but had not been mapped or measured in sufficient detail to demonstrate stratigraphic relations. Our work demonstrates the following stratigraphy (from base to top): (1) Table Mountain Latite (TML) trachyandesite (latite) and basaltic-trachyandesite (shoshonite) lava flows, with phenocrystic clinopyroxene and skeletal plagioclase and groundmass olivine, and minor olivine basalt and basaltic-andesite lava flows; (2) Tollhouse Flat Member of the Eureka Valley Tuff (EVT) welded ignimbrite, with abundant phenocrystic biotite; (3) a single trachydacite lava flow with phenocrystic amphibole, referred to here as the Lava Flow Member of the EVT; (4) By-Day Member of the EVT welded ignimbrite, which

lacks phenocrystic biotite; (5) Upper Member of the EVT, consisting of unwelded ignimbrite; and (6) a single very thick (60 m) aphyric basaltic-trachyandesite (shoshonite) lava flow that we assign to the Dardanelles Formation.

Our work confirms for the first time that the Dardanelles Formation exists as a unit that overlies all members of the EVT, and that its geochemistry is the same as that of the TML in the type section. Our documentation of a trachydacite lava flow within the EVT in the central Sierra Nevada shows that widespread effusive volcanism alternated with explosive volcanism, rather than merely preceding and following it, and that magmas of trachydacitic composition were erupted as lavas as well as pyroclastic flows. The preservation of all formations and members of the Stanislaus Group in the “Cataract” channel suggests that it was an important paleogeomorphic feature.

INTRODUCTION

The Stanislaus Group of the central Sierra Nevada consists of voluminous, widespread high-K volcanic rocks interpreted to record the birth of the Sierra Nevada microplate through transtensional faulting at about 11–9 Ma (Busby and Putirka, 2009). The Stanislaus group consists of high-K lava flows and pyroclastic flows that were likely erupted from the Little Walker center, 8 km east of the present-day Sierran crest, and flowed mainly westward down a paleochannel/paleocanyon system cut into Mesozoic granitic and metamorphic basement, toward the Central Valley (Figure 1; Ransome, 1898; Lindgren, 1911; Slemmons, 1953; Curtis, 1954; Garside et al., 2005; King et al., 2007; Busby et al., 2008a; Gorny et al., 2009; Busby and Putirka, 2009). Despite the fact that these high-K volcanic rocks have been studied for 120 years, their stratigraphy remains poorly understood, partly due to difficulty of access in roadless wilderness areas. For example, Slemmons (1966) named the Bald Peak-Red Peak area of the Sierra Nevada high country as the type section for the Stanislaus Group, but due to its “relative inaccessibility”, Noble et al. (1974) proposed a reference section in the Sweetwater Mountains along Highway 395 (Figure 1). The stratigraphy of these rocks is also poorly understood because they were deposited in a paleo-river canyon system, where erosion took place between eruptions. It is also likely that different lava flows or pyroclastic flows followed different parts of the paleocanyon system.

This paper focuses on the stratigraphy along a segment of the “Cataract paleochannel” of Ransome (1898) and Lindgren (1911). The “Cataract paleochannel” is a paleo-river canyon that roughly coincides with the modern Stanislaus River along much of its length (Figure 1), so it was referred to as the paleo-Stanislaus River by King et al. (2007). Our new geologic mapping of a segment in the upper reach of this paleo-river canyon (Figure 3) contributes to an understanding of the paleo-morphology of the western edge of the Nevada Plano, by showing that all formations and members of the Stanislaus Group are preserved within it (Figure 4). The study area thus preserves the most complete stratigraphic section through the Stanislaus Group that has been recognized within the Sierra Nevada (Figure 5). This complete section records alternating effusive and explosive volcanism, including (from base to top): basaltic to trachyandesitic effusive volcanism (including shoshonite and latite lava flows; Table Mountain Latite); trachydacitic explosive and effusive volcanism (ignimbrite and lava flow members of the Eureka Valley Tuff); and basaltic-trachyandesite effusive volcanism (i.e. shoshonitic; Dardanelles Formation). This section therefore provides a relatively complete and complex record of magmatic and eruptive processes that operated at the Little Walker center, over a period of at least a million years, when the Sierra Nevada microplate began to calve off the western edge of the Nevada Plano.

PREVIOUS STRATIGRAPHIC WORK

High-K volcanic rocks of the central Sierra Nevada were first recognized in 1898 by Ransome, who described latite lava flows (Table Mountain Latite, TML) overlain by “biotite augite latite”, in turn overlain by the Dardanelles Flow (Figure 5A). The “biotite augite latite” was later recognized as welded and unwelded ash-flow tuffs (ignimbrites) by Slemmons (1966), who referred to it as the Eureka Valley Member of the Stanislaus Formation (Figure 5A). Slemmons (1966) grouped all the high-K volcanic rocks (TML, EVT and Dardanelles Flow) into the Stanislaus Formation, and designated the Bald Peak-Red Peak area as the type section, although, a measured section was not made of the type section. Even though Slemmons (1966) reported latite lava flows within the Eureka Valley Member in the type section, their exact location was not identified, and later reports were not made of them in the central Sierra. Noble et al. (1974) identified

three distinct ignimbrites within the Eureka Valley Member and they raised it to formational status, the Eureka Valley Tuff (EVT), subdivided into the Tollhouse Flat, By-Day and Upper members (Figure 5A). This subdivision required elevation of the Stanislaus Formation to group status (Figure 5A; Noble et al., 1974). Because of difficulty of access to the type section, a reference section was proposed to the east (~20 km) of the Sierra Nevada in the Sweetwater Mountains, 6 km east of highway 395 and north of the Little Walker center (Figure 1; Noble et al., 1974).

For the past 35 years, the stratigraphy of the Stanislaus Group in the central Sierra has been portrayed as a section of quartz latite ignimbrite (EVT members) sandwiched between latite lava flows below and above (the TML and the Dardanelles Formation), as shown on Figure 5A (Noble et al., 1974). The reference section does not include the Dardanelles Formation (King et al., 2007), nor had the Dardanelles Formation been found to rest upon any strata younger than the basal member of the EVT (the Tollhouse Flat Member; Figure 5B). Thus, before our study, it had not been proven that the Dardanelles Formation is younger than all of the units of the EVT Formation. When Hagan et al. (2008) discovered a latite lava flow sandwiched between the Tollhouse Flat and Upper members of the EVT, they questioned the existence of the Dardanelles Formation, and suggested it was actually a member of the EVT. In their interpretation, pyroclastic flow eruptions alternated with, but outlasted, the lava flow eruptions. Our study demonstrates that the high-K pulse of magmatism ended with the eruption of an aphyric basaltic-trachyandesite lava flow.

As is common in all volcanic terranes, the volcanic stratigraphy is more complicated in and around the Little Walker eruptive center, due to emplacement of intrusions and aurally restricted eruptive units. According to Priest (1979), the Little Walker center has latite lava flows that lie between the Tollhouse Flat and By-Day members of the EVT, and Brem (1977) reported a latite lava flow in the same stratigraphic position in another locality (the Sweetwater Mountains), calling it the “Latite Flow Member” of the EVT. New magnetostratigraphic work of Pluhar et al. (2009) shows that this member consists of at least two lava flows, because it has normal polarity at one stratigraphic section, and reversed polarity at another stratigraphic section. Moreover, as discussed below, the lava flow we recognized within the EVT is more

silicic than the latite lava flows Pluhar et al. (2009) describe. Therefore, we cannot follow Brem's (1977) nomenclature for the lava flow between the Tollhouse Flat and the By-Day members (Figure 5B). In an effort to alter the nomenclature established by previous literature as little as possible, we refer to this unit as the Lava Flow Member of the Eureka Valley Tuff, dropping the word "Latite" from its name. Our newly-defined Lava Flow Member of the EVT thus consists of one trachydacite lava flow at the type locality (described here; Figure 7) and at least two trachyandesite (i.e. latite) lava flows outside the study area (Pluhar et al., 2009). We introduce this revised nomenclature with the caveat that it will have to be abandoned if more lava flows are found higher in the section (i.e. between the By-Day and Upper members). For example, in the Little Walker center, Priest (1979) reported 150m (500 ft) of biotite quartz latite lava flows below the Upper Member at Fales Hot Springs. Priest (1979) suggested that the By-Day Member may underlie the biotite quartz latite lava flows, but this was based on the presence of float there. Furthermore, it is unclear from the description whether that the biotite quartz latites are lava flows or intrusion(s). In any event, the Fales Hot Springs unit appears to be restricted to the Little Walker center and is therefore not important to the regional stratigraphy. A complete summary of previous stratigraphic work and new results from the region in and around the Little Walker center are given in Pluhar et al. (2009).

We find that the central Sierran paleocanyon fill contains a trachydacite lava flow within the EVT, between the lower two members (i.e. the Tollhouse Flat and By-Day members), and a basaltic-trachyandesite lava flow on top of the Upper Member of the EVT (Figure 5B); we describe both units so they can be recognized elsewhere. The high-K pulse of volcanism initiated with effusive volcanism (i.e. TML), then alternated between effusive and explosive (i.e. EVT) eruptions, and finally ceased with an effusive eruption (i.e. Dardanelles Fm.). We also present new geochemical data on these units, which shows that the units are all similar in being high in K₂O, but show a large range in silica content.

STRUCTURAL AND STRATIGRAPHIC SETTING

Tertiary volcanic and volcanoclastic rocks of the Bald Peak-Red Peak area lie 7 km west of the Sierra Nevada crest at Sonora Pass (Figure 1). They consist of Oligocene

ignimbrites and Miocene andesitic volcanic and volcanoclastic rocks that dip less than a few degrees and are cut by plugs and a series of NNW-striking faults (Figure 3; Slemmons, 1953).

We mapped two west-dipping normal faults, as well as two intervening, east-dipping normal faults that show only minor offset (Figure 3). We call the eastern fault the Red Peak fault (Figure 3); it dips about 70° westward and has 235 m of vertical separation. Two minor fault splays, located < 1 km to the west of the Red Peak fault, dip 70° eastward and display only 35 m of vertical separation; we interpret them as antithetic faults related to the Red Peak fault. The western fault, herein called the Bald Peak fault, dips $\sim 60^\circ$ westward and has 130 m of vertical separation. Faults are not common west of the range crest in the central Sierra Nevada, although many faults are west of the crest in the southern Sierra (Saleeby et al., 2009).

Stanislaus Group strata are largely absent from the footwall block of the Red Peak fault due to erosion of the upthrown side. We speculate that the Bald Peak fault displacement began during the accumulation of the Miocene section, because a fluvial unit within the Table Mountain Latite (Tstmf) is confined to the downthrown block (Figure 3), and because landslide megablocks several meters in length lie within the debris flow deposits of the Relief Peak Formation (Trpdf) adjacent to the fault. Similarly, abrupt thickening of the Tollhouse Flat Member of the EVT onto the downthrown block of the antithetic faults may indicate that it (and, if related, the Red Peak fault) began to move during emplacement of the Stanislaus Group. Map units clearly thicken and thin in places, however, where we see no evidence for faulting (Figure 3), so these thickness changes may simply record evolving paleotopographic effects within the paleocanyon.

The Oligocene and Miocene strata in Figure 3 lie within a much broader paleochannel/paleocanyon, cut into Mesozoic granitic and metamorphic basement (unconformity 1, Figure 2), whose axis was probably centered at Sonora Peak (Figure 1). The oldest paleocanyon fill deposits consist of several Oligocene nonwelded to welded rhyolite ignimbrites, referred to as the Valley Springs Formation (Figure 2; Slemmons, 1953) which erupted from calderas in central Nevada and flowed westward down paleochannels across the present-day Sierra Nevada to the Sacramento Valley of central California (Garside et al., 2005; Henry, 2008). Thus surface elevations must have

continuously decreased in that direction, from the drainage divide of the Nevada Plano in central Nevada westward to central California, and the region could not have yet been significantly disrupted by normal faults (Busby and Putirka, 2009). Oligocene ignimbrites of the central Sierra were deeply eroded in early Miocene time (unconformity 2, Figure 2; Busby et al., 2008a, 2008b; Busby and Putirka, 2009; Hagan et al., 2009), and overlain by middle Miocene andesitic volcanic and volcanoclastic rocks, referred to as Relief Peak Formation in the Sonora Pass region (Figure 2; Slemmons, 1953, 1966). In the eastern part of the map presented here (Figure 3), the Valley Springs Formation (Tvs) is preserved as an erosional remnant on a paleo-ledge in granitic basement (Kgu). Here the Valley Springs Formation consists of light-colored welded ignimbrite with fiamme and crystals of plagioclase, sanidine and quartz. The Valley Springs Formation is overlain by andesitic debris flow deposits of the Miocene Relief Peak Formation (Trpdf) along a steep buttress unconformity (Figure 3).

In general, Miocene andesitic strata pass gradationally westward from primary and vent-proximal volcanic deposits at the present-day Sierran Crest into reworked volcanoclastic deposits exposed along modern west-flowing drainages; this, together with sparse paleocurrent indicators, provides evidence for continued westward flow of material down the paleocanyons in Miocene time (Busby et al., 2008b). In the area mapped in Figure 3, the Relief Peak Formation consists entirely of dark-tan-colored, unsorted, matrix-supported, polymict, coarse pebble to small boulder debris flow deposits composed of heterogeneous clasts of andesitic composition in a sandstone matrix. Basement granitic clasts are rare. Some blocks in the debris flow deposits are prismatically-jointed or retain delicate breadcrust texture on their surfaces, indicating that they were derived from block-and-ash flows and not transported far from the vent area (Busby et al., 2008b). The debris flow deposits were emplaced cold, as indicated by the presence of non-charred petrified wood fragments.

The Relief Peak Formation (as young as 10.39 ± 0.18 Ma; Table 1; Figure 2) is separated from overlying high-K volcanic rocks of the 10.41–9.34 Ma (Table 1; Figure 2) Stanislaus Group by a third erosional unconformity (Figure 2; Busby et al., 2008b). Measurements of elongate vesicles in lava flows from the map area (Figure 3) provide further evidence for westward flow down the “Cataract” paleocanyon (Figure 6). The

Table Mountain Latite is the most widespread unit of the Stanislaus Group, extending from east of the Sierran crest to the Sierra foothills near Knight's Ferry (Figure 1; Gorny et al., 2009). The Eureka Valley Tuff consists of unwelded to densely welded ignimbrite that range in composition from trachydacite to dacite (Figure 7; King et al., 2007). Its basal member, the Tollhouse Flat Member, has been considered to be the most voluminous and widespread of the three ignimbrite members (Noble et al., 1974). Published descriptions of the Dardanelles Formation contradict one another, probably because it has been confused with the unit we call the Lava Flow Member of the EVT (see discussion section below).

The early Late Miocene Stanislaus Group is in turn separated from the overlying Late Miocene andesitic volcanic and volcanoclastic rocks of the Disaster Peak Formation by a fourth unconformity (Figure 2; Busby et al., 2008b; Busby and Putirka, 2009). In the map area of Figure 3, strata of the Disaster Peak Formation are limited to a single small erosional remnant (Tpdf). Additionally, we assign 7.28 ± 0.06 Ma (Table 1; Figure 2) hornblende-plagioclase-phyric andesite plugs that cut the Stanislaus Group to the Disaster Peak Formation, even though they are intrusions, because they clearly represent the subvolcanic parts of it (Table 1; Figure 3).

Several previous workers inferred that the source for the EVT lies in the Little Walker center (Noble et al., 1974; Priest, 1979; see discussion in King et al., 2007). Slemmons (1966), however, portrayed the source of the Table Mountain Latite as a 36 km by 8 km, WNW-trending region of "latite intrusives" that encompasses the Leavitt Peak - Sonora Peak - Bald Peak - Red Peak area (Slemmons, 1953, 1966). His interpretation has been repeated in more recent publications (Noble et al., 1974; King et al., 2007). We have mapped this entire region in detail and find no evidence for the assertion by Slemmons — just a few small hornblende andesite plugs and dikes. The vent area for a lava flow field of the magnitude of the TML would likely be obvious in this very well-exposed area, just as it is for the 16 Ma Lovejoy basalt in the northern Sierra Nevada (Garrison et al., 2008). The fact that latite lava flows are interstratified with EVT in and around the Little Walker center (Priest, 1979) indicates to us that that all of the high-K rocks vented there. The Little Walker center/caldera is interpreted to have formed along a releasing stepover of a dextral transtensional fault zone (Putirka and Busby, 2007;

Busby et al., 2008b; Busby and Putirka, 2009). According to this explanation, the faults of this zone penetrated a thick crustal section into lithospheric mantle, thereby providing a conduit for low-degree, high-K partial melts. We mapped the area in Figure 3, which lies ~10 km to the west of the Little Walker center (Figure 1), to investigate this distinctive high-K pulse of magmatism.

STANISLAUS GROUP STRATIGRAPHY

The purpose of this paper is to document field relations at the type section of the Stanislaus Group, where exposure is nearly 100% (Figures 3, 5).

Table Mountain Latite

The Table Mountain Latite (TML) lava flows are easily recognized in the field by their clinopyroxene and skeletal plagioclase phenocrysts, up to 1 cm in size (Figure 4), accompanied by groundmass olivine. Geochemical analyses from a measured section through the TML at Sonora Peak 9 km to the ESE, presented by Busby et al., (2008b), show that it ranges from trachyandesite to basaltic-trachyandesite (Table 2; Figure 7). The TML is continuous across the map area (Figure 3), because much of the relief on the paleocanyon floor was filled in by the underlying Relief Peak Formation. The unconformity between the Relief Peak Formation and the Stanislaus Group (unconformity 3; Figure 2) has less relief in the map area (Figure 3) than other areas (e.g. Sonora Pass, Busby et al., 2008b). On Red Peak in the east, the TML consists of seven or eight latite flows, 160 m thick with the top eroded. Five latite flows compose the 150 m thick TML section between the two faults (Figure 4) and west of the Bald Peak fault the TML has four latite flows (Figure 3). Thus, the number of flows decreases westward over a very short distance (just a few kilometers). This westward paleoflow direction is consistent with the fact that only 9 kilometers to the ESE on Sonora Peak, the TML section is 405 m thick and contains 26 flows (top eroded; Busby et al., 2008b). This eastward thickening of TML is compatible with an eruptive source in the Little Walker caldera, which is 15 km east of the map area (Figures 1, 3).

We subdivided the TML section in the Bald Peak-Red Peak area on the basis of outcrop characteristics (Figure 4). Future paleomagnetic and geochemical analysis of

these rocks may enable correlation of specific Bald Peak-Red Peak flows over long distances and provide additional age constraints by correlation to the magnetic polarity timescale (for examples see Busby et al., 2008b; Gorny et al., 2009; and Pluhar et al., 2009). Most of the flows in the map area have prominent flow-top breccias, and some exhibit flow-bottom breccias, but some flows lack breccia and probably represent pahoehoe flows (Figure 4). Such variations are also present at Sonora Peak (Busby et al., 2008a, 2008b).

The basal flow in the TML at Bald Peak-Red Peak has the best developed columnar jointing, probably due to quenching upon what was then the base of the paleocanyon, perhaps in water. On Red Peak the basal TML flow is flow 2 rather than TML flow 1, because flow 1 is absent there. TML flow 1 consists of two flows (1 and 1', Figure 4) but neither is continuous across the map area. Flow 1 has more crystals than flow 1' and is marked by a vesicular top in the line of measured section, although the vesicular top passes laterally into a flow-top breccia. Flow 1' has distinctive bombs and plastically-deformed spatter on its upper surface. Flow 2 has larger augite crystals than the underlying and overlying flows. Flow 3 has laterally-continuous contorted flow banding and a continuous flow-top breccia, but laterally variable plagioclase content. Flow 4 has the most abundant plagioclase of all the flows (30–35%) and flow 4' has the largest plagioclase (up to 10mm) and clinopyroxene phenocrysts (up to 7 mm) of all the flows; perhaps this is equivalent to the “large plagioclase member” (Priest, 1979), which is the second member from the base of the TML at the Little Walker center. Flow 4 is restricted to Red Peak and extends east of Figure 3, whereas flow 4' is continuous across the map area. Flow 5 displays a distinctive weathering pattern that yields jagged blocks.

Fluvial deposits intervene between flows 4 and 5 (Tstmf, Figure 4), exemplified by the one channelized body at the line of the measured section. As noted above, the fluvial deposits thicken dramatically onto the downthrown block west of the Bald Peak fault (Figure 3). There, flow 5 lenses out within a minor channel cut into the fluvial deposits. The fluvial deposits consist of well-sorted, subrounded to rounded pebble to boulder conglomerate and pebbly sandstone and lesser sandstone, in thin to thick beds with planar lamination and cut and fill structures. They contain a mixture of andesite and

latite clasts, and rare granitic clasts, with maximum particle sizes ranging from 15 to 200 cm.

Discontinuous, black, olivine-plagioclase-phyric basaltic and basaltic-andesite lava flows lie between flows 3 and 4 (Figure 3). On the west side of the Bald Peak fault, the geochemistry of the single flow there is basaltic, whereas the single, thicker flow on the east side of the fault is a basaltic-andesite (Figure 7), indicating that two different olivine-plagioclase-phyric lava flows are present at this stratigraphic horizon.

Eureka Valley Tuff

Tollhouse Flat Member

The basal member of the EVT, the Tollhouse Flat Member, consists entirely of densely welded ignimbrite in the map area (Figures 3, 4). It is dominated by black vitrophyre and large black fiamme up to 54 cm long with perlitic texture. It is distinguished from the By-Day Member by abundant biotite phenocrysts (in addition to plagioclase, present in both), and it has bigger and more abundant fiamme that weather out to form cavities. In the map area, the Tollhouse Flat Member also contains about 10–20% dominantly pebble-sized accidental rock fragments. The Tollhouse Flat Member is only 7 m thick in the line of the measured section (Figure 3), and lenses out abruptly to the west of it (Figure 4), reappearing abruptly on the west side of the Bald Peak fault where it is about 25 m thick. It also thickens abruptly onto the downthrown block of the antithetic faults, where it maintains an even thickness of about 25 m to the Red Peak fault.

Lava Flow Member

The Lava Flow Member of the EVT, which lies between the Tollhouse Flat and By-Day members, is mapped here for the first time in the Sierra Nevada. However, as mentioned above, a lava flow member has been recognized at this stratigraphic position in the region of the Little Walker center (Priest, 1979) and in the Sweetwater Mountains (Brem, 1977). At Bald Peak - Red Peak, the Lava Flow Member of the EVT is distinguishable from the TML by its smaller phenocrysts and the presence of plagioclase laths, rather than skeletal or resorbed plagioclase. The Lava Flow Member of the EVT at Bald Peak-Red Peak differs from all other units of the Stanislaus Group by having

amphibole phenocrysts. As discussed below, its chemistry is quite distinct from that of all the other lava flows in the Stanislaus Group described to date, because it is a trachydacite, rather than a trachyandesite or basaltic-trachyandesite. It has the same composition as pumice fragments from the Tollhouse Flat Member of the EVT (Figure 7). In the map area of Figure 3, the Lava Flow Member of the EVT has a very-highly-vesiculated top with a distinctive purple or orange color, which passes laterally into a thick flow-top breccia in the line of the measured section (Figure 4). The Lava Flow Member is continuous wherever it is overlain by By-Day Member of EVT, but it is locally cut out along an erosional surface at the base of the Dardanelles Formation, 375 m west of the Bald Peak fault (Figure 3).

By-Day Member

The By-Day Member of the EVT lacks biotite phenocrysts, although it contains biotite in the groundmass, and consists of strongly welded vitrophyric (black) to weakly-welded devitrified (dark gray) ignimbrite. Its accidental lithic content is similar to that of the Tollhouse Flat Member. It locally has an orange top that was probably produced by vapor phase alteration. The By-Day Member thins from 30 m by the Red Peak fault to 20 m in the line of the measured section, where it underlies Upper Member EVT; it is absent west of the Bald Peak fault (Figure 3).

Upper Member

The Upper Member of the EVT is preserved as erosional remnants on the ridge between the Red Peak and Bald Peak faults and consists of up to 8 m of unwelded biotite-bearing ignimbrite with 20–35% white unflattened pumices. It forms a distinctive soft white outcrop. In the line of the measured section, the ignimbrite is underlain by a fine-grained white tuff 20 cm thick that may be an ash fall deposit.

Dardanelles Formation

The Dardanelles Formation consists of a single lava flow up to 60 m thick, underlain by the Upper Member of the EVT or the Lava Flow Member of the EVT, and overlain by andesitic fluvial sandstones and conglomerates of the Disaster Peak Formation (Figures 3, 4). (Figure 3). In thin section the Dardanelles Formation contains microcrystalline olivine, pyroxene, and trachytic-textured plagioclase laths. It has

extremely irregular jointing and a thick pink-red flow-top breccia injected by coherent lava from the flow's interior. Locally, it has a flow-bottom breccia. It is preserved as erosional remnants up to 60 m thick on the ridge between the Red Peak and Bald Peak faults, and is 40 m thick where it is intruded by the Bald Peak plug and where its flow-top breccia is mostly obliterated or eroded.

Stanislaus Group Age Controls

Age controls on the TML include an Ar/Ar age of 10.41 ± 0.08 Ma for the basal flow, and 10.36 ± 0.06 Ma for the top flow, of the 23 flows on Sonora Peak (Figure 2; Busby et al., 2008b). The Sonora Peak section is mainly composed of normal polarity lava flows, but contains two reversed polarity zones, each represented by a single lava flow (Busby et al., 2008b; Pluhar et al., 2009). The lower reversed lava flow exhibits what we call the "Classic Table Mountain Latite" remanence direction, which found widely in the TML, at sites from Sonora Peak to the foothills (see summary in Pluhar et al., 2009). This direction records an interlude that Pluhar et al. (2009) informally call the Table Mountain Event, which occurred between 10.36 ± 0.06 Ma and 10.41 ± 0.08 Ma, around the time of subchron C5n.2n-2 (10.309–10.313 Ma; Evans et al., 2007). In the foothills near Knight's Ferry (Figure 1), the TML consists of at least four flows, with paleomagnetic results consistent with the "Classic Table Mountain Latite" direction and inconsistent with directional results from any other Stanislaus Group lava flows published to date (Gorny et al., 2009). Assuming typical secular variation rates, the lava flows of the "Classic Table Mountain Latite" were emplaced in less than four centuries (Pluhar et al., 2009).

Precise ages for the By-Day and Upper members of the EVT (9.42 ± 0.04 and 9.43 ± 0.02 Ma, respectively; Pluhar et al., 2009) coupled with their normal polarity (King et al., 2007) indicates that these units were emplaced early in subchron C4Ar.1n (9.443–9.351 Ma; Evans et al., 2007).

The Dardanelles Formation has a single whole rock K-Ar date of 9.3 ± 0.4 , on a sample described an aphyric basalt lava flow at an elevation of 9,230' on Bald Peak by Dalrymple (1964). Our mapping (Figure 3) confirms that this is indeed the Dardanelles Formation, although the Dardanelles Formation is a shoshonite, not a basalt, as described

below (Figure 7). Like the underlying By-Day and Upper members of the EVT, the Dardanelles Formation at Bald Peak displays normal polarity (King et al., 2007). Taken together, these observations suggest that Dardanelles Formation erupted during one of two possible time periods: (1) during normal subchron C4Ar.1n (9.443–9.351 Ma), soon after EVT, or (2) during normal chron C4An (9.098–8.769 Ma; Lourens et al., 2004). Ar/Ar dating in progress will likely distinguish between these two possibilities.

PRELIMINARY GEOCHEMICAL RESULTS

Geochemical analyses of the TML samples demonstrate that their compositions range from trachyandesite (latite) to basaltic-trachyandesite (shoshonite; Figure 7). Trachyandesite is equivalent to a latite for these rocks (Figure 7; Ransome, 1898; Slemmons, 1953), but note that a few TML samples range in composition from andesite to basaltic-trachyandesite. These samples might be considered the more mafic equivalents of the latite series and are still related to the latite lava flows of the TML. Basaltic-trachyandesite is equivalent to a shoshonite. The close relation of these data points suggests that these flows are part of the same magmatic event.

The Lava Flow Member of the EVT plots in the same field as the EVT ignimbrites and is markedly more silicic than all the other lava flows of the Stanislaus Group discovered to date (the TML and Dardanelles formations; Figure 7). This trachydacite lava flow member is geochemically closely related to the Tollhouse Flat Member of the EVT (Figure 7), indicating that the lava flow must have erupted from the same magma chamber in which the EVT was developed, during the same magmatic event. Perhaps it records tapping of a lower, relatively volatile-poor part of the same magma chamber that erupted the Tollhouse Flat Member of the EVT, or tapping of the upper part of the chamber after the gas-rich part vented. In the second interpretation, the By-Day Member represents magma erupted when gasses once again collected at the higher parts of the magma chamber.

As described above, Pluhar et al. (2009) have identified a lava flow with normal polarity in one locality and reversed polarity in another locality in the region of the Little Walker caldera, in the same stratigraphic position (between Tollhouse Flat and By-Day members of the EVT) as our newly-defined Lava Flow Member of the EVT. Preliminary

geochemical data on the lava flows identified by Pluhar, however, indicate that they are indeed latites (with compositions ranging from trachyandesite/latite to andesite; Pluhar, pers. comm., 2009). Thus, Pluhar et al. (2009) follow Brem's (1977) nomenclature by referring to the lava flow as the Latite Flow Member of the EVT. In the type section (Figures 3, 4), that member is instead a trachydacite, so we have dropped the term "Latite" from the name of this unit and refer to it as the Lava Flow Member of the EVT. Our new geochemical data thus indicate that at least three lava flows are included the Lava Flow Member the trachydacite lava flow described here, and the two different trachyandesite (latite) lava flows recognized by Pluhar et al. (2009). Future work will determine whether the trachydacite lava flow has normal or reverse magnetization.

The Dardanelles Formation plots near the same geochemical field as the TML, although it is slightly more mafic in composition than most TML flows (Figure 7). The three samples from the Dardanelles Formation differ in total alkali content; one is basaltic trachyandeiste (i.e shoshonite; BP046) and the others are basaltic-andesite (AAK-08-824-6 and AAK-08-909-2; Table 2). However, the sample that is low in total alkalis may have been altered.

As a final observation, regarding volcanic rock nomenclature, our geochemical data at Sonora Pass, the type locality for latite, provide something of a test of geochemical classification schemes. Le Bas et al. (1986), for example, propose that the term "latite" be used to describe trachyandesites that have high K_2O , specifically, where $K_2O > (Na_2O - 2.0)$. In their terminology, latite is a subdivision of trachyandesite, and rocks that have $Na_2O - 2.0 > K_2O$ are termed benmoreite. Lava flows that occur at Sonora Peak span a somewhat broad range of compositions, but all have $K_2O > (Na_2O - 2.0)$, and most are largely consistent with the Le Bas (1986) definition. The Le Bas et al. (1986) definition is in turn largely consistent with the original intent of Ransome (1898), when he originated the term latite to describe the extrusive form of monzonite. These are rocks compositionally intermediate between trachyte and andesite, though Ransome (1898) perhaps did not foresee the subdivision of trachyandesites into high Na and high K suites. In any case, by these definitions, which we judge to be fully appropriate, not all (but most) samples at Sonora Peak within the TML are latites. Four samples (of 21 total analyzed flows) plot within the basaltic-trachyandesite field (Figure 7); these samples,

also having $K_2O > Na_2O - 2.0$, can be termed shoshonites (as opposed to mugearites, which have $K_2O < (Na_2O - 2.0)$; Le Bas et al., 1986). In addition, one flow from Sonora Pass also plots in the basaltic-andesite field, so it is not trachytic. As for nomenclature, our data thus verify the usefulness and accuracy of the classification scheme as outlined by Le Bas et al. (1986), at least for use of the term latite. In addition, our data further show that latites, at their type locality at least, are a product of shoshonite fractionation, and that shoshonites can in turn be derived by fractionation of basaltic-andesite parent magmas.

DISCUSSION

Previous workers misidentified the Lava Flow Member of the EVT as part of the Dardanelles Formation. Slemmons (1966) reported multiple flows in the Dardanelles Formation (Tsd) but at Bald Peak-Red Peak area, but we recognize only one. Moreover, lava flows previously assigned to the Dardanelles Formation directly overlie the Tollhouse Flat Member of the EVT in many localities (Figure 5B). As a result, it is unclear whether some of the lava flows formerly attributed to the Dardanelles Formation are actually part of the Lava Flow Member of the EVT; without the By-Day Member of EVT in between to demarcate them. It is possible that they have been lumped together in previous interpretations. Previous workers (Ransome, 1898; Slemmons, 1966) have described the Dardanelles Formation (Tsd) on the Dardanelles (Figure 1) as a rock that differs from the TML by lacking skeletal plagioclase texture and containing smaller phenocrysts. However, this description of the Dardanelles Formation is similar to our newly-recognized Lava Flow Member of the EVT at the Bald Peak-Red Peak area (Figure 4).

The Bald Peak-Red Peak area, where stratigraphic relationships are clear and stratigraphy is relatively complete, provides an excellent locale for clearly distinguishing between the TML, the Lava Flow Member of the EVT, and the Dardanelles Formation. The petrographic, geochemical, and outcrop characteristics defined here may be used to extend stratigraphic control into adjacent areas. All the lava flows are distinguishable on the basis of geochemical analyses. The TML consists of plagioclase-clinopyroxene-phyric (basaltic-trachyandesite/shoshonite and trachyandesite/latite) lava flows, with minor interstratified basalt and basaltic-andesite lava flows. In our study area, the Lava

Flow Member of the EVT is plagioclase-pyroxene-amphibole-phyric, in contrast to all other previously described lava flows of the Stanislaus group, which lack amphibole. Furthermore, the Lava Flow Member of the EVT has the same composition as the EVT ignimbrites it is interstratified with (trachydacite); this may record nearly coeval tapping of gas-rich and gas-poor parts of the magma chamber, or volatile-phase re-stratification processes within the magma chamber. Further age controls, in the form of paleomagnetic work and high-precision Ar/AR dating, together with detailed geochemical studies may resolve this question. The Dardanelles Formation is distinctive in that it appears to consist of one aphyric lava flow that is much darker in color and more mafic in composition than most flows of the TML or the EVT.

CONCLUSIONS

In this paper we present the first detailed geologic map and the first measured section of the type locality for the Stanislaus Group. These data demonstrate a complex stratigraphy, due to erosional surfaces between formations and members, but also demonstrate that the type locality contains the most complete stratigraphic section of the Stanislaus Group yet recognized in the Sierra Nevada. All formations and members of the Stanislaus Group were guided down the “Cataract” paleocanyon, which must have been an important paleo-morphological feature. This work thus refocuses attention on Slemmons’ type section (1966) and away from Noble et al.’s (1974) more convenient reference section. Our work also recognizes the first known effusive equivalent of the Eureka Valley Tuff (i.e. the Lava Flow Member), with the same trachydacitic composition.

The Stanislaus Group records a distinctive high-K pulse of magmatism. Volcanism alternated between effusive and explosive eruptive styles, and complexly alternated between mafic, intermediate and silicic compositions, over a period of at least 1 my. Future work will determine how many of the outcrop, petrographic, geochemical and paleomagnetic characteristics of lava flows and ignimbrites discussed here can be used for regional-scale correlations

ACKNOWLEDGMENTS

Our research was supported by National Science Foundation grant EAR-01252 (to Busby, Gans and Skilling), EAR-0711276 (to Putirka and Busby), and EAR-0711181 (to Busby). Formal reviews by Arthur Sylvester and David Wagner were very helpful. The detailed map (Figure 3) was built upon unpublished mapping at a larger scale by Dylan Rood (2003), with support from USGS EDMAP grant 03HQAG0030 (to Busby). Field assistance to Koerner from Megan Gambs and Adrienne Kentner (2008) is gratefully acknowledged. We also acknowledge CSUF undergraduate students, Tim Burns, Shanon Lodge, Chastity Mendez, Matthew Owens, and Louis Tesseo for analyzing the EVT samples. We thank the Kennedy Meadows Pack Station for rock hauling. Thanks is extended to Paul Renne for assistance in recalculating previously-published Ar/Ar dates, using new decay constants.

CAPTIONS

Figure 1 - Distribution of high-K volcanic rocks of the central Sierra Nevada (Stanislaus Group), modified from King et al., 2007. Inset shows physiographic setting: B&R – Basin and Range; CR – Coast Ranges; GV – Great Valley; KM – Klamath Mountains; SCM – Southern Cascade Mountains; SN – Sierra Nevada.

Figure 2 –Stratigraphy of volcanic and volcanoclastic rocks in the Sonora Pass region of the central Sierra Nevada, modified from Busby et al. (2008b). Relative thicknesses of units are extremely variable and are not shown to scale. Age data (recalculated from Busby et al., 2008b) are summarized in Table 1. Figure 3 map units are depicted on the right side of the diagram. Formations are divided by lithofacies wherever possible.

Figure 3 – Geologic map of the Bald Peak-Red Peak area, southern Carson-Iceberg Wilderness, central Sierra Nevada, California: mapping by Alice Koerner in 2008 at a scale of 1:6,000. Map location is outlined on Figure 1. Key to map units is displayed on Figure 2. In the TML, individual lava flows are mapped and numbered; these and other

units of the Stanislaus Group are described in detail in the measured section shown in Figure 4.

Figure 4 – Measured section through the Stanislaus Group at Slemmons’ (1966) type section between Bald Peak and Red Peak (line of section plotted on Figure 2), measured by Alice Koerner in 2008. Although Slemmons described all of the formations and members measured here (see text for discussion), a detailed measured section or geologic map of the area were not provided. Plag = plagioclase, cpx = clinopyroxene, ol = olivine, avg. = average, xtal = crystal, EVT = Eureka Valley Tuff.

Figure 5 – (A) – Evolution of the stratigraphic nomenclature of the Stanislaus Group, modified from King et al. (2007) to include the new data reported in this paper. (B) – Columnar sections summarizing the distribution of various units of the Stanislaus Group: ¹ denotes stratigraphy summarized by King et al (2007); ² denotes stratigraphy described by Hagan et al., 2008; and “this study” refers to the stratigraphy mapped and measured in Figures 3 and 4 of this paper.

Figure 6 – Rose diagram of trends of elongate vesicles within lava flows of the Stanislaus Group in the area mapped in Figure 3 (n = 22). The arrow denotes the average of the trends (255°), which is consistent with the general trend of the “Cataract” paleocanyon (Figure 1).

Figure 7 – Le Bas (1986) plot of high-K rocks and basalts of the Stanislaus Group from the Bald Peak-Red Peak and Sonora Pass areas (locations given on Table 2; samples from the Bald Peak-Red Peak area plotted on Figure 3). This plot shows that the Lava Flow Member of the EVT plots in the same field as the ignimbrites of the EVT. Also, the undated Dardanelles Formation, which we show lies at the top of the Stanislaus Group above all members of the EVT, plots in the same field as the distinctive plagioclase-clinopyroxene-phyric flows of the TML.

Table 1 – Age controls on Sonora Pass stratigraphy. Interpreted ages as reported by Gans (Busby et al., 2008b) are recalculated here using improvements to decay constant (Renne et al., 1998; Kuiper et al., 2008), resulting in preferred ages plotted on Figure 2. Sample localities for samples collected and analyzed from the field area (i.e. BP068) are plotted on Figure 3.

Table 2 – Summary of sample localities and rock names for geochemical data plotted in Figure 7. TML = Table Mountain Latite; EVT = Eureka Valley Tuff; Dard = Dardanelles Formation; ¹type section, Figure 4, this paper; ²see Figure 3, this paper; ³reference section, King et al., 2007; ⁴pumices, from Long Barn, CA; ⁵Sonora Peak measured section, Busby et al., 2008b. *FeO was reported by ³King et al., 2007, whereas Fe₂O₃ was reported by this study. All coordinates reported using WGS 84. The samples^{1,2,4,5} shown in this table were generated from replicate analyses with initial totals between 99.5-100.5, and the replicates were averaged and renormalized.

REFERENCES

- Al-Rawi, Y., 1969, Cenozoic history of the northern part of Mono basin, California and Nevada [Ph.D. dissertation], University of California, Berkeley.
- Busby, C., DeOreo, S., Skilling, I., Gans, P, and Hagan, J., 2008a, Carson Pass-Kirkwood paleocanyon system: implications for the Tertiary evolution of the Sierra Nevada, California: Geological Society of America Bulletin, p. 274-299.
- Busby, C.J., Hagan, J., Putirka, K., Pluhar, C., Gans, P., Rood, D., DeOeo, S., Skilling, I. and Wagner, D., 2008b, The ancestral Cascades arc: Implications for the development of the Sierran microplate and tectonic significance of high K₂O volcanism: in, J. Wright and J. Shervais (eds), Ophiolites, Arcs and Batholiths: Geological Society of America Special Paper 438, p. 331-378.

- Busby, C.J., and Putirka, K., 2009, Miocene evolution of the western edge of the Nevada Plano in the central and northern Sierra Nevada: paleocanyons, magmatism, and structure: in, Gary Ernst, editor, *The Rise and Fall of the Nevada Plano*, International Geology Reviews.
- Brem, G. F., 1977, *Petrogenesis of Late Tertiary potassic volcanic rocks in the Sierra Nevada and western Great Basin*: [PhD dissertation], University of California, Riverside.
- Curtis, G.H., 1954, Mode of origin of pyroclastic debris in the Mehrten Formation of the Sierra Nevada: *University of California Publications in Geological Sciences*, v. 29, p. 453-502.
- Dalrymple, G. B., 1964, *Cenozoic chronology of the Sierra Nevada*: *University of California Publications in Geological Sciences*, v. 47, p. 41.
- Evans, H. F., Westerhold, T., Paulsen, H., and Channell, J. E. T., 2007, Astronomical ages for Miocene polarity chrons C4Ar-C5r (9.3–11.2 Ma), and for three excursion chrons within C5n.2n: *Earth and Planetary Science Letters*, v. 256, no. 3-4, p. 455-465.
- Garrison, N., Busby, C.J., Gans, P.B., Putirka, K. and Wagner, D.L., 2008, A mantle plume beneath California? The mid-Miocene Lovejoy flood basalt, northern California: in, J. Wright and J. Shervais (eds), *Ophiolites, Arcs and Batholiths: A Tribute to Cliff Hopson*: *Geological Society of America Special Paper 438*, p. 551-572.
- Garside, L.J., Henry, C.D., Faulds, J.E., and Hinz, N.H., 2005, The upper reaches of the Sierra Nevada auriferous gold channels, in Rhoden, H.N., ed., *Window to the World*, *Geological Society of Nevada Symposium Proceedings*, May 14–18, 2005.

Gorny, C., Busby, C., Pluhar, C., Hagan, J., and Putirka, K., 2009 this volume, An in-depth look at distal Sierra Nevada paleochannel fill: drill cores through the Table Mountain Latite near Knight's Ferry: in, Gary Ernst, editor, The Rise and Fall of the Nevada Plano, International Geology Reviews. 18 ms pp.

Hagan, J.C., Putirka, K., and Busby, C.J., 2008, Revision to the stratigraphy of the high-potassium volcanic rocks of the Stanislaus Group in the central Sierra Nevada: Geological Society of America Abstracts with Programs, v. 40, no. 1, p. 59.

Hagan, J., Busby, C., Renne, P., and Putirka, K., 2009 this volume, Cenozoic evolution of paleochannels, Ancestral Cascades magmas, and range-front faults in the Kirkwood - Carson Pass - Markleeville region (Central Sierra Nevada): in, Gary Ernst, editor, The Rise and Fall of the Nevada Plano, International Geology Reviews.

Henry, C. D., 2008, Ash-flow tuffs and paleovalleys in northeastern Nevada: Implications for Eocene paleogeography and extension in the Sevier hinterland, northern Great Basin: Geosphere, v. 4, p. 1-35.

King, N., Hillhouse, J., Sherman, G., Hausback, B, and Pluhar, C., 2007, Stratigraphy, paleomagnetism, and anisotropy of the Miocene Stanislaus Group, central Sierra Nevada and Sweetwater Mountains, California and Nevada: Geosphere, v. 3; no. 6; p. 646-666.

Le Bas, M.J., Le Maitre, R.W., Streckeisen, A., and Zenettin, B., 1986, A chemical classification of volcanic rocks based on the total alkali-silica diagram: Journal of Petrology, v. 27, p. 745-750.

Lindgren, W., 1911, The Tertiary Gravels of the Sierra Nevada of California: Washington D.C., United States Geological Survey, p. 222.

- Noble, D.C., Slemmons, D.B., Korrington, M.K., Dickinson, W.R., Al-Rawi, Y., and McKee, E.H., 1974, Eureka Valley Tuff, east-central California and adjacent Nevada: *Geology*, v. 2, p. 139-142.
- Pluhar, C., Deino, A., King, N., Busby, C., Hausback, B., Wright, T., and Fischer, C., 2009 this volume, Magnetostratigraphy, geochronology and proposed revision of Stanislaus Group stratigraphy: in, Gary Ernst, editor, *The Rise and Fall of the Nevada Plano*, International Geology Reviews.
- Priest, G.R., 1979, Geology and geochemistry of the Little Walker volcanic center, Mono County, California: [PhD dissertation], Oregon State University.
- Putirka, K., and Busby, C.J., 2007, The tectonic significance of high K₂O volcanism in the Sierra Nevada, California: *Geology*, v. 35, no. 10, p. 923 – 926.
- Ransome, F.L., 1898, Some lava flows of the western slope of the Sierra Nevada, California, U.S.G.S. Survey Bulletin, p. 89.
- Saleeby, J., Saleeby, Z., Nadin, E., and Maheo, G. , 2009, Step over in the structure controlling the regional west tilt of the Sierra Nevada microplate: eastern escarpment system to Kern Canyon system: in, Gary Ernst, editor, *The Rise and Fall of the Nevada Plano*, International Geology Reviews.
- Slemmons, D.B., 1953, Geology of the Sonora Pass region [PhD thesis]: Berkeley, University of California.
- Slemmons, D.B., 1966, Cenozoic volcanism of the central Sierra Nevada, California, geology of northern California: California Division of Mines and Geology Bulletin, v. 170, p. 199-208.




Figure 1

Sonora Pass Stratigraphy





Figure 2

Stanislaus Group

		Graphic Column	Rock type	Color	Unit	Description	Phenocrysts			
Dardanelles Formation	242.0		shoshonitic aphyric lava flow	black	Tsd	flow top breccia	aphyric; ol and pyx in thin section			
	15.2					coherent		flow bottom breccia		
	Eureka Valley Tuff	226.8		Upper Member	white	Tseu	unwelded tuff	unconformity bio, plag		
		8.2					ash layer			
		218.4		By-Day Member	purple	Tseb	weakly welded ignimbrite	plag		
							20		densely welded ignimbrite	
		198.4		Lava Flow Member	light gray or purple	Tsel	scoriaceous, spinose texture	7% plag, euhedral laths 0.2-4 mm, 1-3% pyx, an-subhedral <1mm; oxidized amphibole <1mm		
							35		flow top breccia with scoria bombs	coherent
							163.4		planar flow banding	flow bottom breccia
		Table Mountain Latite	150.4		Tollhouse Flat Member	black	Tset	welded ignimbrite	bio, plag	
			13							
			114.4		latite lava flow	dark gray	Tstml 5	flow top breccia	5% plag, skeletal 8 mm, 4 mm rounded xtals <2% cpx, subhedral 2-3 mm; ol?	
	36							jagged-edged cliff former		platy parting
	106.4			fluvial units	tan	Tstmf	well-stratified and well-sorted			
6										
86.4			latite lava flow	dark gray	Tstml 4'	flow top breccia	20% plag-rich, skeletal laths 5-10 mm, not aligned; 5-7% cpx, euhedral, up to 7 mm			
						22		continuous cliff-former	flow bottom breccia	
61.4			latite lava flow	dark blue-gray	Tstml 3	flow top breccia	7% plag laths and round xtals, 2-5 mm, aligned; 1% cpx, an-subhedral 0.2-2 mm			
						25		coherent, contorted flow banding	flow bottom breccia	
28.4		latite lava flow	dark blue-gray	Tstml 2	elongate vesicles (40%)	20-25% plag skeletal laths and round xtals, avg. 4-7 mm, not aligned; 3-5% cpx, sub-euhedral 2-5 mm				
					33		cliff with shear faces	scoriaceous section	bulbous cliff former	flattened vesicles (20%) 2-20 mm
14.4		latite lava flow	dark gray-green	Tstml 1'	flow top breccia with plastically deformed bombs and scoria	15-20%, plag laths and round xtals 8-10 mm, aligned; <1% cpx, <1 mm				
					14		elongate vesicles			
					14.4		vesiculated (20%) 5-7mm			
0		andesitic debris flow deposits	Brown	Trpdf	columnar jointed, bulbous cliff	unconformity				
					0		polymict andesitic debris flow deposits			

Total Thickness (m)
 Thickness (m)
 section continues downward into Relief Peak Formation

Figure 4

A

Ransome (1898)	Slemmons (1966)		Noble et al. (1974)		Koerner et al. (this study)			
Dardanelle Flow	Stanislaus Group	Dardanelles Member	Stanislaus Group	Dardanelles Formation		Stanislaus Group	Dardanelles Formation	
		Eureka Valley Member		Eureka Valley Tuff	Upper Member		Eureka Valley Tuff	Upper Member
					By-Day Member			By-Day Member
Biotite-Augite Latite					Tollhouse Flat Member			Lava Flow Member
Table Mountain Flow	Table Mountain Latite Member	Table Mountain Latite		Table Mountain Latite				

B

Locations									
Rattlesnake Hill ¹	McKays Point ¹	Whittakers Dardanelles ¹	Bald Peak ¹	Bald Peak (this study)	Bald Peak - Red Peak area (this study)	Golden Canyon ²	Sonora Pass ¹	EVT, Reference Section, Sweetwater Mountains ¹	Bodie and Mono Lake ¹
		Dardanelles Formation	Dardanelles Formation	Dardanelles Formation	Dardanelles Formation		Dardanelles Formation		
					Upper Member, EVT	Upper Member, EVT		Upper Member, EVT	Upper Member, EVT
					By-Day Member, EVT	Latite Flow Member, EVT		By-Day Member, EVT	
				Latite Flow Member, EVT	Latite Flow Member, EVT				
	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT	Tollhouse Flat Member, EVT
Table Mountain Latite	Table Mountain Latite	Table Mountain Latite		Table Mountain Latite	Table Mountain Latite	Table Mountain Latite	Table Mountain Latite	Table Mountain Latite	

Figure 5.




Figure 6.




Figure 7

Sonora Pass Geochronology													
Sample	Geochemistry	Mineral	Lat (°N)	Long (°W)	WMPA	IsoA	Interpreted Age (Ma) ⁰		Nominal Age (Ma) ¹		Preferred Age (Ma) ²		Unit Name
							Age	± 2σ	Age	± 2σ	Age	± 2σ	
BP068	Andesite	Hbl	38.37609	119.76975	7.12±0.06	7.26±0.16	7.12	0.06	7.22	0.06	7.28	0.06	Hbl Andesite Plug - Disaster Peak Fm
BP068	Andesite	Plag	38.37609	119.76975	7.04±0.50	6.83±0.94	7.0	0.5	7.11	0.5	7.15	0.5	Hbl Andesite Plug - Disaster Peak Fm
TF003	-	Plag	38.43096	119.44792	9.11±0.04	9.10±0.08	9.14	0.04	9.28	0.04	9.34	0.04	Upper Member, Eureka Valley Tuff - Stanislaus Gp
TF003	-	Bio	38.43096	119.44793	9.18±0.04	9.20±0.04	9.18	0.04	9.32	0.04	9.38	0.04	Upper Member, Eureka Valley Tuff - Stanislaus Gp
TF005b	-	Plag	38.43041	119.44805	9.19±0.32	9.10±0.52	9.2	0.3	9.34	0.3	9.4	0.3	By-Day Member, Eureka Valley Tuff - Stanislaus Gp
TF009	-	Plag	38.42891	119.44841	9.27±0.04	9.30±0.10	9.27	0.04	9.41	0.04	9.47	0.04	Tollhouse Flat Member, Eureka Valley Tuff - Stanislaus Gp
TF009	-	Bio	38.42891	119.44842	9.35±0.04	9.32±0.06	9.34	0.04	9.48	0.04	9.54	0.04	Tollhouse Flat Member, Eureka Valley Tuff - Stanislaus Gp
PC032	Shoshonite	Plag	38.35378	119.6344	10.14±0.06	10.15±0.08	10.14	0.06	10.30	0.06	10.36	0.06	Uppermost Table Mtn Latite Flow - Stanislaus Gp
PC005	Latite	Plag	38.34641	119.63263	10.19±0.08	10.30±0.16	10.19	0.08	10.35	0.08	10.41	0.08	Lowermost Table Mtn Latite Flow - Stanislaus Gp
PC-BA	Basaltic andesite	Hbl	38.34427	119.6340	10.10±0.06	10.17±0.18	10.17	0.18	10.33	0.18	10.39	0.18	Block-and-ash flow tuff - Upper Relief Peak Fm
PC-BA	Basaltic andesite	Plag	38.34427	119.6341	na	na	~10	na	~10	na	~10	na	Block-and-ash flow tuff - Upper Relief Peak Fm
BP057	-	Plag	38.37824	119.7430	na	na	23.8	0.2	24.16	0.2	24.32	0.2	Uppermost welded ignimbrite - Valley Springs Fm

Notes:

⁰Interpreted age is calculated using 27.60 Ma for the FCs standard as reported by Gans (Busby et al., 2008b)

¹Nominal age is calculated using 28.02 Ma for the FCs standard (Renne et al., 1998)

²Preferred age is calculated using 28.201 Ma for the FCs standard (Kuiper et al., 2008)

WMPA is data reported by Busby et al., 2008b

IsoA is data reported by Busby et al., 2008b

Table 1

Sample	Fm	Map Unit	Geochemistry	Igneous Fm	Lat (°N)	Long (°W)	SiO ₂	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃ *	MnO	MgO	CaO	Na ₂ O	K ₂ O	P ₂ O ₅	Total	Total Alkalis
AAK-08-824-6	Dard ¹	Tsd ¹	Basaltic andesite	Lava	38.37145	119.76038	54.20	1.19	18.28	8.81	0.09	4.16	7.34	2.60	2.68	0.66	99.99	5.28
AAK-08-909-2	Dard ¹	Tsd ¹	Basaltic andesite	Lava	38.37735	119.74738	54.84	1.14	18.44	8.66	0.09	4.32	7.70	2.68	1.75	0.39	100.00	4.43
BP046	Dard ¹	Tsd ¹	Shoshonite	Lava	38.37413	119.75879	53.79	1.22	18.17	7.87	0.09	4.11	7.44	3.70	2.88	0.72	100.00	6.58
LWC 76G	EVT ³	Tseu ³	Dacite	Ash-flow tuff	38.43019	119.44800	65.88	0.91	17.34	3.71*	0.11	2.00	2.73	2.44	4.68	0.19	99.99	7.12
LWC 85	EVT ³	Tseu ³	Trachydacite	Ash-flow tuff	38.42990	119.45144	64.75	0.94	16.94	2.95*	0.08	0.95	2.99	3.52	6.75	0.13	100.00	10.27
LWC 65d	EVT ³	Tseb ³	Trachydacite	Ash-flow tuff	38.42951	119.44802	62.63	1.17	16.94	4.57*	0.09	1.46	3.23	3.73	5.75	0.42	99.99	9.48
LWC 70d	EVT ³	Tseb ³	Trachydacite	Ash-flow tuff	38.42992	119.44804	63.06	1.14	16.87	4.53*	0.09	1.19	3.01	4.31	5.4	0.4	100.00	9.71
LWC 59G	EVT ³	Tseb ³	Trachydacite	Ash-flow tuff	38.42947	119.44800	61.12	1.19	18.00	4.98*	0.10	1.72	3.73	4.22	4.48	0.46	100.00	8.70
AAK-08-825-2	EVT ¹	Tsel ¹	Trachydacite	Lava	38.37442	119.76389	63.46	1.12	17.10	4.93	0.04	1.24	2.94	3.46	5.34	0.37	100.00	8.80
AAK-08-830-2	EVT ¹	Tsel ¹	Trachydacite	Lava	38.37298	119.75229	64.36	1.10	17.20	4.73	0.05	1.06	2.70	3.18	5.30	0.37	100.02	8.48
LWC 83	EVT ³	Tset ³	Trachydacite	Ash-flow tuff	38.42865	119.44839	64.56	0.86	17.10	3.70*	0.07	0.99	2.70	4.53	5.19	0.30	100.00	9.72
LWC 52G	EVT ³	Tset ³	Trachydacite	Ash-flow tuff	38.42798	119.44791	64.92	0.85	17.08	3.55*	0.07	0.93	2.66	4.55	5.11	0.29	100.01	9.66
LWC 56G	EVT ³	Tset ³	Trachydacite	Ash-flow tuff	38.42845	119.44828	64.82	0.90	17.09	3.68*	0.05	0.86	2.62	4.42	5.25	0.30	99.99	9.67
LWC 82	EVT ³	Tset ³	Trachydacite	Ash-flow tuff	38.42785	119.44759	64.71	0.85	17.04	3.63*	0.08	1.08	2.69	4.60	5.04	0.29	100.01	9.64
LWC 54d	EVT ³	Tset ³	Trachydacite	Ash-flow tuff	38.42845	119.44828	64.89	0.87	17.08	3.82*	0.08	0.72	2.59	4.61	5.03	0.30	99.99	9.64
CMEVT-2R NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	66.39	0.79	16.69	3.60	0.12	1.03	2.00	3.71	5.31	0.35	100.00	9.02
CMEVT-3P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.31	1.01	16.95	4.11	0.11	1.34	2.77	4.54	4.56	0.30	100.00	9.10
CMEVT-5R NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.98	0.91	17.49	3.84	0.09	1.27	2.64	3.67	4.60	0.51	100.00	8.27
LTEVT-P1 NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	65.29	0.91	16.80	3.59	0.10	1.11	2.47	4.62	4.86	0.25	100.00	9.48
LTEVT-P2 NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	65.08	0.94	17.07	4.27	0.10	1.31	2.75	3.36	4.69	0.44	100.00	8.04
LTEVT-P4 NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.96	0.93	17.23	3.77	0.10	1.17	2.51	4.36	4.72	0.25	100.00	9.08
MOEVT-2P 1NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	63.90	0.99	16.98	3.99	0.10	1.31	2.88	5.05	4.50	0.30	100.00	9.56
MOEVT-4P 1NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	66.48	0.73	16.70	3.10	0.10	0.83	1.85	4.74	5.29	0.18	100.00	10.03
MOEVT-3P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.96	1.00	17.12	4.25	0.09	1.31	2.78	3.65	4.54	0.30	100.00	8.19
SLEVT-1P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.80	0.93	17.05	3.56	0.09	1.18	2.61	4.73	4.79	0.26	100.00	9.52
SLEVT-4P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	66.06	0.91	17.01	3.42	0.07	1.13	2.46	3.73	4.97	0.25	100.00	8.70
SLEVT-5P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	66.42	0.84	16.77	3.42	0.10	0.95	2.16	3.80	5.33	0.22	100.00	9.13
TBEVT-1P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.22	1.00	16.84	5.57	0.11	1.42	2.90	3.24	4.38	0.32	100.00	7.62
TBEVT-2P NB	EVT ⁴	Tset ⁴	Dacite	Ash-flow tuff	38.34080	119.74430	64.95	1.08	18.62	4.59	0.10	1.21	2.68	2.79	3.72	0.25	100.00	6.52
TBEVT-3P NB	EVT ⁴	Tset ⁴	Trachydacite	Ash-flow tuff	38.34080	119.74430	64.87	0.96	17.03	3.94	0.10	1.20	2.73	4.16	4.73	0.28	100.00	8.89
SO-P-0107	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	56.30	1.31	18.36	6.40	0.13	2.05	7.78	3.67	3.34	0.67	100.00	7.00
SO-P-0201	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.27	1.31	17.48	7.47	0.05	2.31	6.95	3.44	3.10	0.62	100.00	6.54
SO-P-0305	TML ⁵	Tstm ⁵	Shoshonite	Lava	38.32160	119.68980	55.91	1.26	16.59	7.16	0.08	4.89	7.39	3.10	3.03	0.59	100.00	6.14
SO-P-0402	TML ⁵	Tstm ⁵	Shoshonite	Lava	38.32160	119.68980	55.88	1.25	16.62	6.86	0.11	4.43	8.15	3.14	2.97	0.59	100.00	6.11
SO-P-0505	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.11	1.30	17.09	5.96	0.12	2.36	9.34	3.24	2.88	0.61	100.00	6.12
SO-P-0602	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	55.92	1.52	18.02	6.36	0.16	1.48	8.63	3.60	3.40	0.91	100.00	7.00
SO-P-0708A	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.18	1.57	18.93	6.28	0.02	1.52	6.17	3.80	3.59	0.93	100.00	7.39
SO-P-0804B	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	56.14	1.21	16.72	6.87	0.10	3.68	8.95	2.78	3.03	0.53	100.00	5.80
SO-P-0902	TML ⁵	Tstm ⁵	Shoshonite	Lava	38.32160	119.68980	55.34	1.15	16.57	6.92	0.14	5.41	7.85	3.08	2.98	0.56	100.00	6.06
SO-P-1001	TML ⁵	Tstm ⁵	Basaltic andesite	Lava	38.32160	119.68980	55.47	1.32	16.32	7.65	0.10	5.70	7.39	2.54	2.89	0.61	100.00	5.43
SO-P-1105	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	56.35	1.75	17.72	7.37	0.05	2.44	6.33	3.38	3.62	0.99	100.00	7.00
SO-P-1201	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	56.33	1.75	17.64	7.33	0.07	2.65	6.03	3.51	3.60	1.08	100.00	7.10
SO-P-1304	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.64	1.70	17.92	6.75	0.07	1.44	5.41	3.83	4.26	0.97	100.00	8.09
SO-P-1404	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.63	1.41	18.08	6.54	0.05	2.66	5.91	3.48	3.54	0.71	100.00	7.02
SO-P-1501	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	58.37	1.29	21.16	4.19	0.11	0.80	6.12	3.96	3.40	0.59	100.00	7.36
SO-P-1606	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	58.18	1.42	18.83	6.22	0.08	1.65	5.19	3.51	4.23	0.69	100.00	7.74
SO-P-1806	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	59.16	1.51	18.40	6.42	0.02	1.27	4.53	3.64	4.37	0.69	100.00	8.01
SO-P-1903A	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.15	1.27	17.20	7.51	0.34	2.61	6.48	3.43	3.45	0.55	100.00	6.88
SO-P-2106A	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	58.14	1.67	19.04	5.64	0.01	1.15	5.38	3.75	4.41	0.81	100.00	8.16
SO-P-2204A	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	57.52	1.19	17.69	5.99	0.15	3.58	6.32	3.32	3.70	0.55	100.00	7.02
SO-P-2301	TML ⁵	Tstm ⁵	Latite	Lava	38.32160	119.68980	59.17	1.38	18.32	6.26	0.02	1.30	5.20	3.57	4.19	0.60	100.00	7.76
BP007	TML ²	Tstmb ²	Basalt	Lava	38.36367	119.77765	50.40	1.65	14.94	8.54	0.10	10.15	9.18	2.65	1.91	0.47	100.00	4.57
BP020	TML ²	Tstmb ²	Basaltic andesite	Lava	38.37160	119.77208	56.53	0.89	16.55	6.39	0.07	6.61	7.39	3.70	1.62	0.25	100.00	5.32

Table 2